

Decoupling of Translations from Homogeneous Transformations in Inhomogeneous Quantum Groups*

Gaetano Fiore,

Dip. di Matematica e Applicazioni, Fac. di Ingegneria
Università di Napoli, V. Claudio 21, 80125 Napoli

I.N.F.N., Sezione di Napoli,
Complesso MSA, V. Cintia, 80126 Napoli

Abstract

We briefly report on our result that, if there exists a realization of a Hopf algebra H in a H -module algebra $\mathcal{A}$, then their cross-product is equal to the product of $\mathcal{A}$ itself with a subalgebra isomorphic to H and *commuting* with $\mathcal{A}$. We illustrate its application to the Euclidean quantum groups in $N \geq 3$ dimensions.

1 Introduction and main results

Let H be a Hopf algebra and $\mathcal{A}$ a unital H -module algebra. Here for the sake of brevity we stick to consider the case of a right H -module algebra. Then by definition there exists a bilinear map $\triangleleft : (a, g) \in \mathcal{A} \times H \rightarrow a \triangleleft g \in \mathcal{A}$, called a right action, such that

$$a \triangleleft (gg') = (a \triangleleft g) \triangleleft g' \quad (1)$$

$$(aa') \triangleleft g = (a \triangleleft g_{(1)}) (a' \triangleleft g_{(2)}). \quad (2)$$

*Talk given at the 2nd International Symposium on “Quantum Theory and Symmetries”, July 2001, Krakow, Poland. Preprint 00-45 Dip. Matematica e Applicazioni, Università di Napoli; Preprint DSF/36-2001.

We use a Sweedler-type notation with suppressed summation sign for the coproduct $\Delta(g)$ of g , namely the short-hand notation $\Delta(g) = g_{(1)} \otimes g_{(2)}$. The cross-product algebra $\mathcal{A} \rtimes H$ is $H \otimes \mathcal{A}$ as a vector space, whereas the product of two elements is given by

$$(g \otimes a)(g' \otimes a') = gg'_{(1)} \otimes (a \triangleleft g'_{(2)})a' \quad (3)$$

for any $a, a' \in \mathcal{A}$, $g, g' \in H$. To simplify the notation, in the sequel we denote $g \otimes a$ by ga and omit either unit $\mathbf{1}_{\mathcal{A}}, \mathbf{1}_H$ whenever multiplied by non-unit elements; consequently, for $g = \mathbf{1}_H$, $a' = \mathbf{1}_{\mathcal{A}}$ (3) takes the form

$$ag' = g'_{(1)} (a \triangleleft g'_{(2)}). \quad (4)$$

$\mathcal{A} \rtimes H$ itself is a H -module algebra under the right action $\triangleleft$ of H if we extend the latter on the elements of H as the adjoint action,

$$h \triangleleft g = Sg_{(1)}hg_{(2)}, \quad g, h \in H. \quad (5)$$

One very important situation in which one encounters cross products is in inhomogeneous (quantum) groups, that are Hopf algebras whose algebras are cross-products $\mathcal{A} \rtimes H$, where H is the Hopf subalgebra of “homogeneous transformations” and $\mathcal{A}$ the (braided) Hopf subalgebra of “translations”.

In Ref. [3] we have found a sufficient condition for $\mathcal{A} \rtimes H$ to be equal to a product $\mathcal{A}H'$, where $H' \subset H \ltimes \mathcal{A}$ is a subalgebra isomorphic to H and *commuting* with $\mathcal{A}$: it is sufficient to assume that there exists an algebra map

$$\tilde{\varphi} : \mathcal{A} \rtimes H \rightarrow \mathcal{A} \quad (6)$$

acting as the identity on the subalgebra $\mathcal{A}$, namely for any $a \in \mathcal{A}$

$$\tilde{\varphi}(a) = a. \quad (7)$$

Here we briefly recall this result, as well as a similar one requiring a weaker assumption, namely that H admits a Gauss decomposition into two Hopf subalgebras H^+, H^- for each of which analogous maps $\tilde{\varphi}^+, \tilde{\varphi}^-$ exist. In the last section we illustrate the application of these general results to a class of inhomogeneous quantum groups, the Euclidean quantum groups in $N \geq 3$ dimensions, for which $\mathcal{A}$ is (“the algebra of functions on”) the quantum Euclidean N -dimensional space; the corresponding maps $\tilde{\varphi}, \tilde{\varphi}^+, \tilde{\varphi}^-$ have been determined in [1] and further studied in [4]. [Note that no such maps can exist for the (undeformed) Euclidean algebra, for which $\mathcal{A} \equiv$ the algebra of functions

on $\mathbf{R}^N$, which is abelian]. Other applications include $U_q\mathfrak{g}$ -covariant Heisenberg (or Clifford) algebras and the q -deformed 2-dimensional fuzzy sphere $S_{q,M}^2$ [5].

Let $\tilde{\mathcal{C}}$ be the commutant of $\mathcal{A}$ within $\mathcal{A} \rtimes H$, i.e. the subalgebra

$$\tilde{\mathcal{C}} := \{c \in \mathcal{A} \rtimes H \mid [c, a] = 0 \quad \forall a \in \mathcal{A}\}. \quad (8)$$

Clearly $\tilde{\mathcal{C}}$ contains the center $\mathcal{Z}(\mathcal{A})$ of $\mathcal{A}$. Let $\tilde{\zeta} : H \rightarrow \mathcal{A} \rtimes H$ be the map defined by

$$\tilde{\zeta}(g) := g_{(1)} \tilde{\varphi}(Sg_{(2)}). \quad (9)$$

Note that if we apply $\tilde{\varphi}$ to $\tilde{\zeta}$ and recall that $\tilde{\varphi}$ is both a homomorphism and idempotent we find $\tilde{\varphi} \circ \tilde{\zeta} = \varepsilon$. Now, $\varepsilon(g)$ is a complex number times $\mathbf{1}_{\mathcal{A}}$ and therefore trivially commutes with $\mathcal{A}$. Since by definition the commutation relations between $\mathcal{A}$ and either H or $\tilde{\varphi}(H)$ are the same, we expect that also $\tilde{\zeta}(g)$ commutes with $\mathcal{A}$. This is confirmed by

Theorem 1 [3] *Under the above assumptions the map $\tilde{\zeta}$ is an injective algebra homomorphism $\tilde{\zeta} : H \rightarrow \tilde{\mathcal{C}}$; moreover $\tilde{\mathcal{C}} = \tilde{\zeta}(H) \mathcal{Z}(\mathcal{A})$ and $\mathcal{A} \rtimes H = \tilde{\zeta}(H) \mathcal{A}$. If, in particular $\mathcal{Z}(\mathcal{A}) = \mathbf{C}$, then $\tilde{\mathcal{C}} = \tilde{\zeta}(H)$ and $\tilde{\zeta} : H \hookrightarrow \tilde{\mathcal{C}}$ is an algebra isomorphism. Moreover, the center of the cross-product $\mathcal{A} \rtimes H$ is given by $\mathcal{Z}(\mathcal{A} \rtimes H) = \mathcal{Z}(\mathcal{A}) \tilde{\zeta}(\mathcal{Z}(H))$. Finally, if $H_c, \mathcal{A}_c$ are maximal abelian subalgebras of H and $\mathcal{A}$ respectively, then $\mathcal{A}_c \tilde{\zeta}(H_c)$ is a maximal abelian subalgebra of $\mathcal{A} \rtimes H$.*

In other words, the subalgebra H' looked for will be obtained by setting $H' := \tilde{\zeta}(H)$. For these reasons we shall call $\tilde{\zeta}$, as well as the other maps $\tilde{\zeta}^{\pm}$ which we shall introduce below, *decoupling maps*.

To introduce the latter we just need a weaker assumption, namely that, instead of a map $\tilde{\varphi}$, we just have at our disposal two homomorphisms $\tilde{\varphi}^+, \tilde{\varphi}^-$

$$\tilde{\varphi}^{\pm} : H^{\pm} \rtimes \mathcal{A} \rightarrow \mathcal{A} \quad (10)$$

fulfilling (7), where H^+, H^- denote two Hopf subalgebras of H such that Gauss decompositions $H = H^+ H^- = H^- H^+$ hold. (The typical case is when $H = U_q\mathfrak{g}$ and H^+, H^- denote its positive and negative Borel subalgebras.) Then Theorem 1 will apply separately to $\mathcal{A} \rtimes H^+$ and $\mathcal{A} \rtimes H^-$, if we define corresponding maps $\tilde{\zeta}^{\pm} : H^{\pm} \rightarrow \mathcal{A}$ by

$$\tilde{\zeta}^{\pm}(g) := g_{(1)} \tilde{\varphi}^{\pm}(Sg_{(2)}), \quad (11)$$

where $g \in H^{\pm}$ respectively. What can we say about the whole $H \rtimes \mathcal{A}$?

Theorem 2 [3] *Under the above assumptions formulae (11) define injective algebra homomorphisms $\tilde{\zeta}^{\pm} : H^{\pm} \rightarrow \tilde{\mathcal{C}}$. Moreover,*

$$\tilde{\mathcal{C}} = \tilde{\zeta}^{+}(H^{+}) \tilde{\zeta}^{-}(H^{-}) \mathcal{Z}(\mathcal{A}) = \tilde{\zeta}^{-}(H^{-}) \tilde{\zeta}^{+}(H^{+}) \mathcal{Z}(\mathcal{A}) \quad (12)$$

and

$$\mathcal{A} \rtimes H = \tilde{\zeta}^{+}(H^{+}) \tilde{\zeta}^{-}(H^{-}) \mathcal{A} = \tilde{\zeta}^{-}(H^{-}) \tilde{\zeta}^{+}(H^{+}) \mathcal{A}. \quad (13)$$

In particular, if $\mathcal{Z}(\mathcal{A}) = \mathbf{C}$, then the commutant is equal to $\tilde{\mathcal{C}} = \tilde{\zeta}^{+}(H^{+}) \tilde{\zeta}^{-}(H^{-}) = \tilde{\zeta}^{-}(H^{-}) \tilde{\zeta}^{+}(H^{+})$. Furthermore, any element c of the center $\mathcal{Z}(\mathcal{A} \rtimes H)$ of the cross-product $\mathcal{A} \rtimes H$ can be expressed in the form

$$c = \tilde{\zeta}^{+} \left(c^{(1)} \right) \tilde{\zeta}^{-} \left(c^{(2)} \right) c^{(3)}, \quad (14)$$

where $c^{(1)} \otimes c^{(2)} \otimes c^{(3)} \in H^{+} \otimes H^{-} \otimes \mathcal{Z}(\mathcal{A})$ and $c^{(1)} c^{(2)} \otimes c^{(3)} \in \mathcal{Z}(H) \otimes \mathcal{Z}(\mathcal{A})$; vice versa any such object c is an element of $\mathcal{Z}(\mathcal{A} \rtimes H)$. Finally, if $H_c \subset H^{+} \cap H^{-}$ and $\mathcal{A}_c$ are maximal abelian subalgebras of H and $\mathcal{A}$ respectively, then $\mathcal{A}_c \tilde{\zeta}^{+}(H_c)$ (as well as $\mathcal{A}_c \tilde{\zeta}^{-}(H_c)$) is a maximal abelian subalgebra of $\mathcal{A} \rtimes H$.

As a consequence of this theorem, for any $g^{+} \in H^{+}$, $g^{-} \in H^{-}$ there exists a sum $c^{(1)} \otimes c^{(2)} \otimes c^{(3)} \in \mathcal{Z}(\mathcal{A}) \otimes H^{-} \otimes H^{+}$ (depending on g^{+}, g^{-}) such that

$$\tilde{\zeta}^{+}(g^{+}) \tilde{\zeta}^{-}(g^{-}) = c^{(1)} \tilde{\zeta}^{-}(c^{(2)}) \tilde{\zeta}^{+}(c^{(3)}). \quad (15)$$

These will be the commutation relations between elements of $\tilde{\zeta}^{+}(H^{+})$ and $\tilde{\zeta}^{-}(H^{-})$. Their form will depend on the specific algebras considered.

Assume that H is a Hopf $*$ -algebra and $\mathcal{A}$ a H -module $*$ -algebra. Then, as known, these two $*$ -structures can be glued in a unique one to make $\mathcal{A} \rtimes H$ a $*$ -algebra itself. What can we say about the behaviour of the decoupling maps under the latter $*$ -structure?

Proposition 1 [3] *If $\tilde{\varphi} : \mathcal{A} \rtimes H \rightarrow \mathcal{A}$ is a $*$ -homomorphism, then also the map $\tilde{\zeta} : H \rightarrow \tilde{\mathcal{C}}$ is. If $\tilde{\varphi}^{\pm}$ are $*$ -homomorphism, then also the map $\tilde{\zeta}^{\pm} : H^{\pm} \rightarrow \tilde{\mathcal{C}}$ are. If $\tilde{\varphi}^{\pm}$ fulfill the condition $\tilde{\varphi}^{\pm}(\alpha^{*}) = [\tilde{\varphi}^{\mp}(\alpha)]^{*}$ for any $\alpha \in H^{\mp} \rtimes \mathcal{A}$, then $\tilde{\zeta}^{\pm}$ fulfill*

$$\tilde{\zeta}^{\pm}(g^{*}) = [\tilde{\zeta}^{\mp}(g)]^{*}, \quad g \in H^{\mp}. \quad (16)$$

2 Application to the Euclidean quantum group $\mathbf{R}_q^N \bowtie U_q so(N)$

As an algebra $\mathcal{A}$ we shall consider a slight extension of the quantum Euclidean space $\mathbf{R}_q^N$ [2] (the $U_q so(N)$ -covariant quantum space), i.e. of the unital associative algebra generated by p^i fulfilling the relations

$$\mathcal{P}_{a_{hk}}^{ij} p^h p^k = 0, \quad (17)$$

where $\mathcal{P}_a$ denotes the q -deformed antisymmetric projector (29), and the indices take the values $i = -n, \dots, -1, 0, 1, \dots, n$ for N odd, and $i = -n, \dots, -1, 1, \dots, n$ for N even; here $n := \left\lfloor \frac{N}{2} \right\rfloor$ is the rank of $so(N)$. The multiplet (p^i) carries the fundamental N -dim (or vector) representation ρ of $U_q so(N)$: for any $g \in U_q so(N)$

$$p^i \triangleleft g = \rho_j^i(g) p^j. \quad (18)$$

As a set of generators of $H \equiv U_q so(N)$ it is convenient to introduce the FRT generators [2] $\mathcal{L}_j^{+i}, \mathcal{L}_j^{-i}$, together with the square roots of the diagonal elements $\mathcal{L}_j^{+j}, \mathcal{L}_j^{-j}$. The FRT generators and the braid matrix $\hat{R}$ are related to the so-called universal R -matrix $\mathcal{R} \in H \otimes H$ by

$$\mathcal{L}_l^{+a} := \mathcal{R}^{(1)} \rho_l^a(\mathcal{R}^{(2)}) \quad \mathcal{L}_l^{-a} := \rho_l^a(\mathcal{R}^{-1(1)}) \mathcal{R}^{-1(2)} \quad (19)$$

and $\hat{R}_{hk}^{ij} = \rho_h^j(\mathcal{L}_k^{+i}) = (\rho_h^j \otimes \rho_k^i)(\mathcal{R})$. Since in our conventions $\mathcal{R} \in H^+ \otimes H^-$ (H^+, H^- denote the positive, negative Borel subalgebras) we see that $\mathcal{L}_l^{+a} \in H^+$ and $\mathcal{L}_l^{-a} \in H^-$.

To define $\tilde{\varphi}$ or $\tilde{\varphi}^\pm$ one [1] slightly enlarges $\mathbf{R}_q^N$ as follows. One introduces some new generators $\sqrt{P_a}$, with $1 \leq a \leq \frac{N}{2}$, together with their inverses $(\sqrt{P_a})^{-1}$, requiring that

$$P_a^2 = \sum_{h=-a}^a p^h p_h = \sum_{h,k=-a}^a g_{hk} p^h p^k. \quad (20)$$

In the previous equation g_{hk} denotes the ‘metric matrix’ of $SO_q(N)$:

$$g_{ij} = g^{ij} = q^{-\rho_i} \delta_{i,-j}. \quad (21)$$

It is a $SO_q(N)$ -isotropic tensor and is a deformation of the ordinary Euclidean metric. We have introduced the notation [2] $(\rho_i) = (n -$

$\frac{1}{2}, \dots, \frac{1}{2}, 0, -\frac{1}{2}, \dots, \frac{1}{2} - n$) for N odd, $(n-1, \dots, 0, 0, \dots, 1-n)$ for N even. In the sequel we shall call P_n^2 also P^2 . Moreover for odd N we add also $\sqrt{p^0}$ and its inverse as new generators. The commutation relations involving these new generators can be fixed consistently, in particular one finds that

$$\sqrt{P_a} p^i = p^i \sqrt{P_a} \times \begin{cases} 1 & \text{if } |i| \leq a \\ \sqrt{q} & \text{if } i > a \\ 1/\sqrt{q} & \text{if } i < -a \end{cases} \quad (22)$$

The center of $\mathbf{R}_q^N$ is generated by $\sqrt{P}$ and, only in the case of even N , by $\sqrt{p^1/p^{-1}}$ and its inverse $\sqrt{p^{-1}/p^1}$. In the case of even N one needs to include also the FRT generator $\mathcal{L}^{-1}_1 = \mathcal{L}^{+1}_{-1}$ and its inverse $\mathcal{L}^{+1}_1 = \mathcal{L}^{-1}_{-1}$ [which are generators of $U_q so(N)$ belonging to the natural Cartan subalgebra] among the generators of $\mathcal{A}$. They satisfy nontrivial commutation relations with the generators of $\mathcal{A}$, and the standard FRT relations with the rest of $U_q so(N)$. As a consequence, $\sqrt{p^{\pm 1}/p^{\mp 1}}$ are eliminated from the center of $\mathcal{A}$ (in fact they do not q -commute with $\mathcal{L}^{\pm 1}_1$).

The homomorphisms $[1] \tilde{\varphi}^{\pm} : \mathcal{A} \rtimes U_q^{\pm} so(N) \rightarrow \mathcal{A}$ take the simplest and most compact expression on the FRT generators of $U_q^{\pm} so(N)$. Let us introduce the short-hand notation $[A, B]_x = AB - xBA$. The images of $\tilde{\varphi}^{\pm}$ on the FRT generators read

$$\tilde{\varphi}^{-}(\mathcal{L}^{-i}_j) = g^{ih} [\mu_h, p^k]_q g_{kj}, \quad (23)$$

$$\tilde{\varphi}^{+}(\mathcal{L}^{+i}_j) = g^{ih} [\bar{\mu}_h, p^k]_{q^{-1}} g_{kj}, \quad (24)$$

where

$$\begin{aligned} \mu_0 &= \gamma_0 (p^0)^{-1} & \bar{\mu}_0 &= \bar{\gamma}_0 (p^0)^{-1} & \text{for } N \text{ odd,} \\ \mu_{\pm 1} &= \gamma_{\pm 1} (p^{\pm 1})^{-1} \mathcal{L}^{\pm 1}_1 & \bar{\mu}_{\pm 1} &= \bar{\gamma}_{\pm 1} (p^{\pm 1})^{-1} \mathcal{L}^{\mp 1}_1 & \text{for } N \text{ even,} \\ \mu_a &= \gamma_a P_{|a|}^{-1} P_{|a|-1}^{-1} p^{-a} & \bar{\mu}_a &= \bar{\gamma}_a P_{|a|}^{-1} P_{|a|-1}^{-1} p^{-a} & \text{otherwise,} \end{aligned} \quad (25)$$

and $\gamma_a, \bar{\gamma}_a \in \mathbf{C}$ are normalization constants fulfilling the conditions

$$\begin{aligned} \gamma_0 &= -q^{-\frac{1}{2}} h^{-1} & \bar{\gamma}_0 &= q^{\frac{1}{2}} h^{-1} & \text{for } N \text{ odd,} \\ \gamma_{\pm 1} &= -k^{-1} & \bar{\gamma}_{\pm 1} &= k^{-1} & \text{for } N \text{ even,} \\ \gamma_1 \gamma_{-1} &= -q^{-1} h^{-2} & \bar{\gamma}_1 \bar{\gamma}_{-1} &= -q h^{-2} & \text{for } N \text{ odd,} \\ \gamma_a \gamma_{-a} &= -q^{-1} k^{-2} \omega_a \omega_{a-1} & \bar{\gamma}_a \bar{\gamma}_{-a} &= -q k^{-2} \omega_a \omega_{a-1} & \text{for } a > 1. \end{aligned} \quad (26)$$

Here $h := q^{\frac{1}{2}} - q^{-\frac{1}{2}}$, $k := q - q^{-1}$, $\omega_a := (q^{\rho_a} + q^{-\rho_a})$. One can choose the γ_a 's, $\bar{\gamma}_a$'s: 1) only for odd N , in such a way that $\tilde{\varphi}^-, \tilde{\varphi}^+$ can be “glued” into a unique homomorphism φ of the type (6), (7); 2) for $|q| = 1$ or $q \in \mathbf{R}^+$ respectively, in such a way that either condition necessary for the application of Proposition 1 is fulfilled. From definition (9), using (30), (23), (24), we find

$$\begin{aligned}\tilde{\zeta}^-(\mathcal{L}_j^{-i}) &= \mathcal{L}_h^{-i} \tilde{\varphi}^-(S\mathcal{L}_j^{-h}) = \mathcal{L}_h^{-i} [\mu_h, p^i]_q, \\ \tilde{\zeta}^+(\mathcal{L}_j^{+i}) &= \mathcal{L}_h^{+i} \tilde{\varphi}^+(S\mathcal{L}_j^{+h}) = \mathcal{L}_h^{+i} [\bar{\mu}_h, p^i]_{q^{-1}}.\end{aligned}\quad (27)$$

In Ref. [3] we have determined the commutation relations (15) between them.

Appendix

The braid matrix of $U_q so(N)$ is given by

$$\begin{aligned}\hat{R} &= q \sum_{i \neq 0} e_i^i \otimes e_i^i + \sum_{\substack{i \neq j, -j \\ \text{or } i=j=0}} e_i^j \otimes e_j^i + q^{-1} \sum_{i \neq 0} e_i^{-i} \otimes e_{-i}^i \\ &+ k \left(\sum_{i < j} e_i^i \otimes e_j^j - \sum_{i < j} q^{-\rho_i + \rho_j} e_i^{-j} \otimes e_{-i}^j \right).\end{aligned}\quad (28)$$

It admits the orthogonal projector decomposition

$$\hat{R} = q\mathcal{P}_s - q^{-1}\mathcal{P}_a + q^{1-N}\mathcal{P}_t; \quad (29)$$

$\mathcal{P}_a, \mathcal{P}_t, \mathcal{P}_s$ are q -deformed antisymmetric, trace, trace-free symmetric projectors. The coproduct and antipode of the FRT generators are given by

$$\Delta(\mathcal{L}_j^{\pm i}) = \mathcal{L}_h^{\pm i} \otimes \mathcal{L}_j^{\pm h}, \quad S\mathcal{L}_i^{\mp j} = g_{ih} \mathcal{L}_k^{\mp h} g^{kj}. \quad (30)$$

References

- [1] B. L. Cerchiai, G. Fiore, J. Madore, “ Geometrical Tools for Quantum Euclidean Spaces”, *Commun. Math. Phys.* **217** (2001), 521-554.
- [2] L.D. Faddeev, N.Y. Reshetikhin, L. Takhtadjan, *Alge. i Anal.* **1** (1989), 178, translated from the Russian in *Leningrad Math. J.* **1** (1990), 193.

- [3] G. Fiore, *On the Decoupling of the Homogeneous and Inhomogeneous Parts in Inhomogeneous Quantum Groups*, Preprint 00-31 DMA, Università di Napoli, DSF/3-2001 and math.QA/0101218.
- [4] G. Fiore, H. Steinacker and J. Wess “Unbraiding the braided tensor product”, math/0007174.
- [5] H. Grosse, J. Madore, H. Steinacker, “Field Theory on the q -deformed Fuzzy Sphere”, hep-th/0005273.